




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/13, 15/62, C07K 14/705, A61K 48/00, 38/17		A1	(11) International Publication Number: WO 99/64597
			(43) International Publication Date: 16 December 1999 (16.12.99)
(21) International Application Number: PCT/US99/12309 (22) International Filing Date: 3 June 1999 (03.06.99) (30) Priority Data: 60/088,813 10 June 1998 (10.06.98) US (71) Applicant (for all designated States except US): THE GOVERNMENT OF THE UNITED STATES OF AMERICA, represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; National Institutes of Health, Office of Technology Transfer, Suite #325, 6011 Executive Boulevard, Rockville, MD 20852-3804 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): RIBAUDO, Randall, K. [US/US]; 11800 Dewey Road, Silver Spring, MD 20906-4881 (US). SHIELDS, Michael [US/US]; 8507 Capricorn Way #80, San Diego, CA 92126 (US). (74) Agent: NOONAN, William, D.; Klarquist, Sparkman, Campbell, Leigh & Whinston, LLP, Suite 1600, One World Trade Center, 121 S.W. Salmon Street, Portland, OR 97204 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: β_2 MICROGLOBULIN FUSION PROTEINS AND HIGH AFFINITY VARIANTS

(57) Abstract

β_2 -microglobulin fusion proteins and modified forms of β_2 -microglobulin are disclosed. The fusion proteins are shown to incorporate onto the surface of MHC I expressing mammalian cells and to cause an increased cytotoxic T-cell response to antigens presented by such cells. The fusion proteins are useful in methods of tumor therapy. Modified forms of human β_2 -microglobulin, particularly a form having a serine to valine transition at amino acid position 55 of the mature protein are shown to have an enhanced affinity for MHC I heavy chain, and are useful both in the disclosed fusion proteins and as a vaccine adjuvant where enhanced cytotoxic T-cell response is desired.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

β_2 MICROGLOBULIN FUSION PROTEINS AND HIGH AFFINITY VARIANTS

Field of the Invention

This invention relates to compositions based on β_2 microglobulin, and the use of such compositions in immunological methods pertaining to the targeting of proteins to cell surfaces. The disclosed compositions and methods have particular application to vaccination and tumor therapy.

Background of the Invention

MHC I and activation of cytotoxic T-cells

The beta-2 microglobulin (β_2m) protein is a component of the class I major histocompatibility complex (MHC I). MHC I is formed by the association of β_2m and an alpha protein (also known as the "heavy" chain), which comprises three domains, a1, a2 and a3. MHC I is found on the surface of most types of nucleated cells, where it presents antigens derived from proteins synthesized in the cytosol to CD8⁺ T-cells. Two signals are required for activation of naive CD8⁺ T-cells. The first signal is provided by the interaction of the T-cell receptor (TCR) with the MHC I-antigen complex on the antigen-presenting cell surface. The second signal is generated by the interaction of a ligand on the antigen-presenting cell (APC) with a second receptor present on the T-cell surface. This second signal is termed co-stimulation, and the APC ligand is often referred to as a co-stimulatory molecule. The best characterized co-stimulatory molecules on APCs are the structurally related glycoproteins B7.1 (CD80) and B7.2 (CD86) which interact with the CD28 receptor on the T-cell surface. Activation of CD8⁺ T-cells by these two signals leads to the proliferation of antigen-specific cytotoxic T-cells, which recognize and destroy cells presenting the signaling antigen. These cytotoxic T-cells play an important role in the immunological defense against intracellular pathogens such as viruses, as well as tumors. A detailed presentation of the immunological basis of the cytotoxic T-cell response can be found in Janeway and Travers (*Immunobiology: the immune system in health and disease*, Current Biology Ltd./Garland Publishing, Inc. New York, 1997).

The failure of an exogenous (non-self) antigen to stimulate a cytotoxic T-cell response can result from a block in the above-described cytotoxic T-cell activation pathway at one of many points (see Ploegh, 1998, *Science* 280:248-53). Failure of the cytotoxic T-cell activation pathway is of great significance in two particular areas of medicine: vaccination and tumor immunology.

Cytotoxic T-cells and vaccination

Vaccine technology has focused in recent years on sub-unit vaccines. Sub-unit vaccines comprise isolated pathogen components, such as viral capsid or envelopes, or synthetic peptides that mimic an antigenic determinant of a pathogen-related protein. For example, U.S. Patent No. 4,974,168 describes leukemia associated immunogens that are peptides based on envelope proteins of

a leukemia-associated virus. However, while sub-unit vaccines can stimulate CD4⁺ helper T-cells (which play a key role in humoral immunity), attempts to stimulate CD8⁺ cytotoxic T-cells *in vivo* with such vaccines have been largely unsuccessful. It has been postulated that the reason for this is the inability of the exogenously administered vaccine peptide to associate with the MHC I molecules on the cell surface (Liu, 1997, *Proc. Natl. Acad. Sci. USA* 94:10496-8). In other words, the block in the cytotoxic T-cell activation pathway occurs at the stage where the antigen is loaded into the MHC I molecule.

One proposed solution to this problem is to combine the antigenic peptide with a molecule that is readily taken up into cells (reviewed by Liu, 1997, *Proc. Natl. Acad. Sci. USA* 94:10496-8). Thus, this strategy is based on getting the antigen into the cytosol so that it can join the normal pathway by which antigens are processed for presentation by MHC I. In contrast, Rock et al. (*J. Immunol.* 150:1244-52, 1993) adopted a strategy of enhancing the binding of the vaccine peptide to MHC I already present on the cell surface. Rock et al. (*J. Immunol.* 150:1244-52, 1993) report that the administration of exogenous purified β_2m along with the vaccine peptide produces enhanced loading of the peptide onto MHC I *in vivo* and thereby stimulates a cytotoxic T-cell response against the peptide. The use of exogenous β_2m as a vaccine adjuvant is also described in U.S. Patent No. 5,733,550 (to Rock et al.), which is incorporated herein by reference.

Tumor cells and immune system evasion

Tumor cell immunity is primarily cell-mediated, involving both CD8⁺ cytotoxic T-cells and CD4⁺ helper T-cells. However, despite the fact that tumor cells express tumor-specific proteins that are not recognized as self-antigens by the immune system, they often escape recognition by the immune system. A number of factors may contribute to the ability of tumor cells to evade immune recognition, including the down-regulation of expression of co-stimulatory proteins. TCR stimulation in the absence of co-stimulatory molecules can result in failure to activate the T-cell and the induction of clonal anergy. Thus, down-regulation of co-stimulatory proteins in tumor cells prevents normal activation of T-cells that do bind to tumor antigens on the cell surface, permitting the tumor cell to escape recognition.

Several research groups have attempted to address this issue by removing tumor cells from a patient, providing exogenous co-stimulatory molecules on the surface of the removed tumor cells and then reintroducing the tumor cells to the patient so that immune recognition can occur. For example, European patent application number 96302009.4 describes a method by which tumor cells are removed from a patient, transfected to express both B7 and CD2 (a co-receptor involved in T-cell adhesion and activation) on the tumor cell surface, and then reintroduced to the patient. The reintroduced cells are reported to stimulate a broad immunological response against both the reintroduced transfected tumor cells and the non-transfected tumor cells within the patient's body, resulting in tumor regression.

Adopting an alternative approach to this problem, Gerstmayer et al. (*J. Immunol.* 158:4584-90, 1997) describe a chimeric B7-antibody protein, in which the antibody is specific for the *erbB2* proto-oncogene product. This chimeric molecule localizes specifically on the surface of *erbB2* expressing tumor cells, and presents the B7 co-stimulatory molecule to cytotoxic T-cells, resulting in enhanced proliferation of cytotoxic T-cells. Gerstmayer et al. (*J. Immunol.* 158:4584-90, 1997) thus propose that fusion proteins comprising an anti-tumor antibody and a co-stimulatory molecule could be useful as tumor immunotherapeutics. However, this approach would require prior knowledge and characterization of tumor-specific antigens expressed on the tumor cells of each individual patient, and the use of an antibody specific for that particular type of tumor cell.

Summary of the Invention

The present invention employs various forms of beta-2 microglobulin to address the problems associated with failure of the cytotoxic T-cell activation pathway in both vaccination and tumor therapy. The invention also provides compositions and methods based on β_2m that are broadly applicable to achieve expression of any desired target protein on the surface of any mammalian cell.

In one embodiment, the invention provides fusion proteins comprising a first amino acid sequence and a second amino acid sequence, wherein the second amino acid sequence is a β_2m microglobulin. In particular applications, the first amino acid sequence may be a co-stimulatory protein, such as B7.1 or B7.2, or another protein having immunological activity, such as a cytokine, an integrin or a cellular adhesion molecule. Examples of such proteins include interleukins (e.g., IL-2, IL-12), granulocyte-macrophage colony-stimulating factor (GM-CSF), lymphocyte function-associated proteins (e.g., LFA-1, LFA-3) and intercellular adhesion molecules (e.g., ICAM-1, ICAM-2). In other embodiments, the first amino acid sequence of the fusion protein may be any protein that is desired to be expressed on the surface of a cell. It is shown that these fusion proteins are an effective way to target a desired protein, such as B7, to the outer membrane of a cell. ("B7" is used generically to refer to either B7.1 or B7.2).

With respect to tumor therapy, it is shown that expressing on the surface of a tumor cell a fusion protein comprising a β_2m joined to a co-stimulatory protein can significantly increase the immune response of an animal to the tumor cell. In one example, a fusion protein comprising h β_2m joined to the co-stimulatory protein B7 (and termed B7- β_2m) is targeted to the surface of tumor cells, such that the tumor cells present the B7- β_2m fusion protein to T-cells. These cells are then attenuated and introduced into mice. T-cells removed from these mice were shown to be significantly more active against the same type of tumor cells than equivalent cells from mice treated with tumor cells presenting β_2m only.

The β_2m fusion proteins provided by the invention have wide application in that they are useful to target any desired protein to the outer membrane of a cell. These fusion proteins may be targeted to the surface of a cell in a number of ways. In one approach, cells that express MHC I are

simply incubated with a preparation of the fusion protein, resulting in the incorporation of the fusion protein onto the cell surface. Alternatively, the fusion protein may be introduced into the cell so that it is incorporated into the MHC I pathway. In another approach, a nucleic acid molecule encoding the fusion protein is introduced into a cell by transformation. Expression of this nucleic acid molecule results in the fusion protein being produced within the cell and exported to the cell membrane. Where the fusion protein is to be introduced into the cytosol for export to the outer membrane, or where it will be expressed by a nucleic acid molecule within the cell, it is desirable to include a signal peptide at the N-terminus of the fusion peptide so that the fusion protein is transported to the outer membrane of the cell. The β_2m signal peptide may be used for this purpose. In all of these approaches, the result is that the β_2m fusion protein is presented on the surface of the cell.

In one embodiment, the invention includes nucleic acid molecules encoding the disclosed fusion proteins, as well as nucleic acid vectors comprising such nucleic acid molecules. Transgenic cells comprising these nucleic acid molecules are also provided by the invention.

Methods of expressing a β_2m fusion protein on the surface of a cell are provided by the invention. Such methods include contacting a cell with a fusion protein comprising a first amino acid sequence and a second amino acid sequence wherein the second amino acid sequence is a β_2m . An alternative method provided by the invention comprises transforming the cell with a nucleic acid molecule encoding such a fusion protein.

The invention further provides methods of enhancing the immune response of an animal to an antigen presented on the surface of a cell. Such methods comprise providing a β_2m fusion protein on the surface of the cell and administering the cell to the animal. In such applications, the fusion protein preferably comprises β_2m fused to a co-stimulatory protein, such as B7, or another protein having immunological activity. Expressing the β_2m fusion protein on the surface of the cell may be accomplished by contacting the cell with the fusion protein, or transforming the cell with a nucleic acid molecule encoding the protein. These methods may be applied to the treatment of tumors; in such treatments, the antigen against which an enhanced immune response is desired is a tumor antigen, and the cell bearing the antigen is a tumor cell. The tumor cell may be removed from the body of a mammal having a tumor, or may be derived from an *in vitro* propagated tumor cell line.

The β_2m fusion protein is introduced to the tumor cell (e.g., by incubation of the tumor cell with the protein, or by transformation of the tumor cell with a nucleic acid encoding the fusion protein), such that the tumor cell presents the fusion protein on its surface. The tumor cell carrying the fusion protein is then administered to a mammal. In certain embodiments, the tumor cell may be attenuated prior to being administered to the mammal; such attenuation may be accomplished using standard means such as radiation, heat or chemical treatment. Once in the body of the mammal, the combination of tumor antigens and the β_2m -fusion protein on the surface of the tumor cells are recognized by CD8⁺ T-cells, resulting in T-cell activation, proliferation and thereby an enhanced

cytolytic T-cell response against both the introduced tumor cells and other tumor cells in the mammal that express the same tumor antigen.

The present invention also provides modified human β_2m ($h\beta_2m$) proteins having an enhanced affinity for MHC I. Such proteins are shown to bind to the alpha chain of MHC I with
5 higher affinity than wild-type $h\beta_2m$ and to enhance T-cell recognition of APCs bearing the modified $h\beta_2m$. In particular embodiments, the modified $h\beta_2m$ proteins have a valine residue at position 55 in place of the serine residue that is found in the mature form of naturally occurring (i.e., wild-type) $h\beta_2m$. Such modified $h\beta_2m$ proteins are referred to as $h\beta_2m$ S55V.

$h\beta_2m$ S55V is useful as a vaccine adjuvant in place of wild-type $h\beta_2m$. Thus, one aspect of
10 the invention is a vaccine preparation comprising at least one antigen and $h\beta_2m$ S55V. $h\beta_2m$ S55V may also be utilized in place of wild-type $h\beta_2m$ in the fusion proteins discussed above. Additionally, β_2m fusion proteins may also be employed in such vaccine preparations, either using a wild-type β_2m or, in the case of $h\beta_2m$, $h\beta_2m$ S55V.

15

Sequence Listing

The nucleic and amino acid sequences listed in the Sequence Listing are shown using standard letter abbreviations for nucleotide bases, and three letter code for amino acids. Only one strand of each nucleic acid sequence is shown, but the complementary strand is understood to be
20 included by any reference to the displayed strand.

Seq. I. D. No. 1 shows the amino acid sequence of wild-type (naturally occurring) $h\beta_2m$.

Seq. I. D. No. 2 shows the amino acid sequence of the B7- β_2m fusion protein (comprising the B7.2 co-stimulatory molecule).

Seq. I. D. No. 3 shows the amino acid sequence of the B7- β_2m fusion protein having the β_2m
25 signal sequence joined to the N-terminal of the B7 domain.

Seq. I. D. Nos. 4-7 show primers used to construct $h\beta_2m$ S55V.

Seq. I. D. Nos. 8 and 9 show primers used to amplify the B7.2 protein.

Seq. I. D. No. 10 shows the amino acid sequence of mature $h\beta_2m$ S55V.

Seq. I.D. Nos. 11 and 12 show the amino acid linker sequences that can be used between the
30 two domains of a fusion protien.

Seq. I.D. Nos. 13 and 14 show amino acid sequences of signal peptides that can be used to direct the expression of a protein in a cell.

Seq. I.D. No. 15 shows the amino acid sequences for a c-myc tag.

Seq. I.D. Nos. 16-20 show the amino acid sequences for peptides used in the HLA
35 stabilization assay.

Brief Description of the Drawings

FIG. 1 shows the amino acid sequence of naturally occurring (i.e., wild-type) $h\beta_2m$; the signal peptide is double underlined and the amino acid numbering starts at the isoleucine residue that is the first residue of the mature protein. The serine at position 55 that is changed to valine in S55V is shown in bold.

FIG. 2 is a schematic representation of the B7.2- β_2m microglobulin fusion peptide construct.

FIG. 3 shows the sequence of the B7.2- β_2m fusion peptide. Residue 1 is a methionine required for expression, residues 2-220 are the extracellular portion of murine B7-2, residues 221-225 (italics) are a sequence created by the insertion of a restriction site into the nucleic acid sequence, residues 226-240 (underlined) are the linker sequence, and residues 241-339 are the mature form of $h\beta_2m$.

FIG. 4 shows the sequence of a B7- β_2m fusion peptide having the $h\beta_2m$ signal sequence (residues 1-20, double underlined). Residues 21-239 are the extracellular portion of murine B7-2, residues 240-244 (italics) are a sequence created by the insertion of a restriction site into the nucleic acid sequence, residues 245-259 (underlined) are the linker sequence, and residues 260-358 are the mature form of $h\beta_2m$.

FIG. 5 is a graph showing the effect of plate-bound B7- β_2m on BALB/c splenic T-cell proliferation in the presence of a suboptimal concentration of soluble 2C11.

FIG. 6 is a graph showing the efficacy of P815 antigen-primed T-cells (primed with P815 cells presenting only P815 antigens, or P815 antigens and either B7- β_2m or β_2m) to lyse P815 tumor cells.

FIG. 7 is a graph illustrating the stabilization of cell-surface HLA-A1 (a), -A2 (b), and -A3 (c) by mutant $h\beta_2m$ and peptide. All values are expressed as mean fluorescence intensity.

FIG. 8 shows the inhibition of myc- β_2m binding by S55V and $h\beta_2m$ to cell-surface HLA-A1 (a), -A2 (b), and -A3 (c). All values are expressed as mean fluorescence intensity.

FIG. 9 illustrates that the S55V mutant enhances CTL recognition better than wild-type $h\beta_2m$ in both Hmy2.C1R-A2 (a) and Hmy2.C1R-A3 (b) target cells.

Detailed Description of the Invention

Abbreviations and Definitions

To facilitate review and understanding of the invention as described herein, the following explanation of abbreviations and definitions of terms are provided:

β_2m : beta-2 microglobulin. This term encompasses any mammalian beta-2 microglobulin protein, including human and murine beta-2 microglobulins. The term " $h\beta_2m$ " refers specifically to human beta-2 microglobulin. cDNAs and genes encoding mammalian β_2m s are well known in the art, as are the corresponding β_2m protein sequences. Examples include those sequences described in: Parnes and Seidman (*Cell* 29:661-9, 1982), Gates et al. (*Proc. Natl. Acad. Sci. USA* 78:554-8, 1981)

(murine); Suggs et al. (*Proc. Natl. Acad. Sci. USA* 78:6613-7, 1981), Guessow et al. (*J. Immunol.* 139:3132-8, 1987), Cunningham et al. (*Biochem.* 12:4811-22, 1973) (human); and Ellis et al. (*Immunogenetics* 38:310, 1993) (bovine). These sequences are also available on GenBank at <http://www.ncbi.nlm.nih.gov/Entrez/index.html>.

5 The terms "wild-type β_2m " and "naturally occurring β_2m " refer to the β_2m protein that is isolated from the particular species of mammal in question. For example, wild-type $h\beta_2m$ refers to a beta-2 microglobulin protein having an amino acid sequence of β_2m isolated from a human source (e.g., serum). Thus, an example of a wild-type β_2m is the $h\beta_2m$ protein disclosed in Cunningham et al. (*Biochem.* 12:4811-4822, 1973), which is also available on GenBank under accession number
10 A90371 and is shown in FIG. 1 and Seq. I.D. No. 1. The term "modified β_2m " refers to a beta-2 microglobulin protein having an amino acid sequence that has been modified from a wild-type β_2m amino acid sequence. By way of example, $h\beta_2m$ S55V is a mutant form of $h\beta_2m$ in which the serine residue present at position 55 in mature wild-type $h\beta_2m$ (see FIG. 1 and Seq. I.D. No. 1) is replaced with a valine residue. The term $h\beta_2m$ S55V encompasses forms of $h\beta_2m$ that differ from wild-type
15 $h\beta_2m$ by the substitution of the position 55 serine for a valine, as well as forms of $h\beta_2m$ that have the S55V modification and additional amino acid sequence modifications.

Fusion protein: A protein comprising two amino acid sequences that are not found joined together in nature. The term " β_2m fusion protein" refers to a protein that comprises a first amino acid sequence and a second amino acid sequence, wherein the second amino acid sequence is a β_2 -
20 microglobulin. The β_2m amino acid sequence and the first amino acid sequence may alternatively be referred to as domains of the fusion protein. Thus, for example, the present invention provides fusion proteins comprising first and second domains, wherein the second domain is a β_2m protein. The link between the first and second domains of the fusion protein is typically, but not necessarily, a peptide linkage. In particular β_2m fusion proteins, the two domains may be joined by means of a
25 linker peptide. In β_2m fusion proteins, the first domain is preferably, but not necessarily, linked to the N-terminus of the β_2m domain.

These fusion proteins may also be represented by the formula X-Y wherein X is a protein, such as a co-stimulatory protein, and Y is a β_2m protein. In a further embodiment of the fusion proteins disclosed, a signal peptide sequence may be linked to the N-terminus of the first protein.
30 Such a three part protein can thus be represented as S-X-Y wherein S is the signal peptide, X is a protein, such as a co-stimulatory protein and Y is a β_2m protein. Where the fusion protein is being expressed in a eukaryotic cell, the signal peptide is preferably a eukaryotic signal peptide that functions to target expression of the fusion protein to the cell membrane. While a number of signal peptides may be used for this purpose, the preferred signal peptide is the β_2m signal peptide (shown
35 in FIG. 1). Where the fusion protein is being expressed in a prokaryotic cell, the signal peptide is preferably a prokaryotic signal peptide that results in the secretion of the fusion peptide into the growth medium from where it can be readily harvested and purified. Suitable prokaryotic signal

peptides are well known in the art. Where the X protein and β_2m are joined by a peptide linker, the fusion protein may be represented as X-L-Y or, if a signal peptide is present, S-X-L-Y, where L is the linker peptide.

Certain β_2m fusion proteins of the present invention include, as their β_2m component, the h β_2m S55V protein. Particular residues in the β_2m component of such fusion proteins may be referred to by the number of residues that they are away from the first residue of the mature h β_2m (which is isoleucine). Thus, the β_2m component of a fusion protein that includes h β_2m S55V may be referred to as a human β_2 -microglobulin having a valine at position 55.

Transformed: A transformed cell is a cell into which has been introduced a nucleic acid molecule by molecular biology techniques. As used herein, the term transformation encompasses all techniques by which a nucleic acid molecule might be introduced into such a cell, including transfection with viral vectors, transformation with plasmid vectors, and introduction of naked DNA by electroporation, lipofection, and particle gun acceleration.

Isolated: An "isolated" biological component (such as a nucleic acid or protein) has been substantially separated or purified away from other biological components in the cell of the organism in which the component naturally occurs, i.e., other chromosomal and extrachromosomal DNA and RNA, and proteins. Nucleic acids and proteins which have been "isolated" thus include nucleic acids and proteins purified by standard purification methods. The term also embraces nucleic acids and proteins prepared by recombinant expression in a host cell as well as chemically synthesized nucleic acids.

Vector: A nucleic acid molecule as introduced into a host cell, thereby producing a transformed host cell. A vector may include nucleic acid sequences that permit it to replicate in the host cell, such as an origin of replication. A vector may also include one or more selectable marker genes and other genetic elements known in the art.

Purified: The term purified does not require absolute purity; rather, it is intended as a relative term. Thus, for example, a purified β_2m protein preparation is one in which the β_2m protein is more pure than the protein in its natural environment within a cell. Preferably, a preparation of a protein is purified such that the protein represents at least 50% of the total protein content of the preparation.

Operably linked: A first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences are contiguous and, where necessary to join two protein coding regions, in the same reading frame.

Recombinant: A recombinant nucleic acid is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated

segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques.

Tumor Cell: A neoplastic cell that may be either malignant or non-malignant. Tumor
5 cells include cells from both solid and non-solid tumors (such as hematologic malignancies). Tumors may be primary tumors originating in a particular organ (such as breast, prostate, bladder or lung). Tumors of the same tissue type may be divided into tumors of different sub-types (a classic example being bronchogenic carcinomas (lung tumors) which can be an adenocarcinoma, small cell, squamous cell, or large cell tumor).

10 **Mammal:** This term includes both human and non-human mammals. Similarly, the term “patient” includes both human and veterinary subjects.

Production and use of β_2m fusion proteins

Standard molecular biology, biochemistry and immunology methods are used in the present
15 invention unless otherwise described. Such standard methods are described in Sambrook et al. (*Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, New York, 1989), Ausubel et al. (*Current Protocols in Molecular Biology*, Greene Publishing Associates and Wiley-Intersciences, 1987), Innis et al. (*PCR Protocols, A Guide to Methods and Applications*, Innis et al. (eds.), Academic Press, Inc., San Diego, California, 1990) and Harlow and Lane (*Antibodies, A Laboratory
20 Manual*, Cold Spring Harbor Laboratory, New York, 1988). Methods of producing nucleic acid sequences expressing fusion proteins, as well as methods of expressing and purifying fusion proteins are well known in the art, and are described, for example, in U.S. patent Nos. 5,580,756 and 5,698,679. The following paragraphs are provided by way of guidance.

Standard techniques may be employed to make genetic constructs expressing β_2m fusion
25 proteins, including restriction endonuclease digestion, ligation and the polymerase chain reaction. Any mammalian gene or cDNA encoding β_2m may be used as the source of the β_2m coding sequence. Such sequences are known in the art and are available on public databases such as GenBank. By way of example, the sequence of the human β_2m cDNA is described in Guessow et al. (*J. Immunol.* 139:3132-8, 1987, GenBank accession number: M17986). Notably, this cDNA
30 sequence includes regions coding for the signal peptide of h β_2m (see FIG. 1).

Similarly, nucleic acid sequences coding for proteins that may be selected as the second domain of the fusion protein are well known in the art. While the selection of the second domain protein will typically be a co-stimulatory protein or a protein having some other immunological activity, fusion proteins may be constructed using β_2m as the second domain, and any protein that is
35 desired to be delivered to the surface of a cell as the first domain. Examples of cDNAs encoding proteins having co-stimulatory activity include those encoding human B7.1 (Freeman et al., 1989, *J. Immunol.* 143:2714-22, GenBank accession number: M27533), B7.2 (Azuma et al., 1993, *Nature*

366:76-9, GenBank accession number L25259), LFA-3 (Wallner et al., 1987, *J. Exp. Med.* 166:923-32, GenBank accession number Y00636) and ICAM-1 (Simmons et al., 1988, *Nature* 331:624-7, GenBank accession number X06990). Examples of other immunologically active proteins include, ICAM-3 (Fawcett et al., 1992, *Nature* 360:481-4, GenBank accession number S50015), VCAM-1
5 (Damle et al., 1992, *J. Immunol.* 148:1985-92), CD59 (Menu et al., 1994, *J. Immunol.* 153:2444-56), CD40 (Hancock et al., 1996, *Proc. Natl. Acad. Sci. USA* 93:13967-72) and GM-CSF (Takashi et al., JP 1991155798-A, GenBank accession number E02975). Proteins having other activities, such as tumor necrosis factor (TNF, Masaaki et al., JP 1985185799, GenBank accession number E00423) may also be employed as the first domain in the fusion protein. By way of example, proteins that
10 induce apoptosis (such as TNF) or anergy may be employed to delete certain classes of antigen-specific activated T-cells. Thus, myelin basic protein (MBP)-specific autoreactive T-cells that are found at the site of inflammation in multiple sclerosis patients may be deleted by introducing into the patient target cells that express MBP and which also present a TNF- β_2 m fusion protein.

cDNA clones encoding β_2 m and the second protein may be obtained as described in the
15 cited references, or by PCR amplification from mRNA (or cDNA libraries) of cells that express the particular protein. cDNA amplification is performed as described by Innis et al. (*PCR Protocols, A Guide to Methods and Applications*, Innis et al. (eds.), Academic Press, Inc., San Diego, California, 1990) using primers designed to amplify the desired portions of the cDNA. For example, cDNA primers may be designed to amplify only that portion of the β_2 m cDNA that encodes the mature form
20 of β_2 m. PCR may also be used to adapt the amplified fragments for ligation.

In addition to a β_2 m domain and a second protein domain (e.g., B7), β_2 m fusion proteins may also include additional elements, such as a linker sequence between the β_2 m domain and the second domain, and a signal peptide. The linker sequence is generally between 10 and 25 amino acids in length, and serves to provide rotational freedom in the fusion construct, thereby facilitating
25 appropriate conformational folding of the two adjacent protein domains. Such linker sequences are well known in the art and include the glycine(4)-serine spacer (GGGGS x3, Seq. I.D. No. 11) described by Chaudhary et al. (*Nature* 339:394-397, 1989). A version of this linker in which the third repeat of the linker motif is modified to GGGAS (Seq. I.D. No. 12) is also shown in FIG. 3 and Seq. I.D. No. 2. Other linker sequences may also be used to construct the β_2 m fusion proteins.

30 Signal peptides serve to direct expression of a particular protein to a specified location in the cell. Depending on whether the fusion protein is to be expressed in a prokaryotic or eukaryotic cell, a prokaryotic or eukaryotic signal peptide will be selected. Prokaryotic signal peptides that direct secretion of peptides into the medium may be particularly useful where large amounts of the fusion peptide are to be produced. Examples of such signal sequences include the prokaryotic signal
35 sequence of the pectate lyase gene pelB (Power et al., 1992, *Gene* 113:95-9 (KYLLPTAAAGLLLLAAQPAMA, Seq. I.D. No. 13)), and the outer membrane protein ompT (Ouzzine et al., 1994, *FEBS Lett* 339:195-9 (MRAKLLGIVLTPIAISFAST, Seq. I.D. No. 14)).

Eukaryotic signal peptides that direct expression of a peptide to the cell surface are useful where the fusion protein is to be presented on the surface of the cell. The signal peptide of β_2m (shown in FIG. 3 and Seq. I.D. No. 2) is particularly suitable for this purpose.

In their most basic form, nucleic acids encoding β_2m fusion proteins will comprise x-y
5 wherein x is a nucleic acid sequence encoding the first protein domain (e.g., B7) and y is a nucleic acid sequence encoding the β_2m protein domain. Where a linker sequence is to be included, the nucleic acid may be represented as x-l-y, wherein l is a nucleic acid sequence encoding the linker peptide. Where a signal sequence is to be included, the nucleic acid may be represented as s-x-l-y wherein s is a nucleic acid sequence encoding the signal peptide. Preferably, although not
10 necessarily, the relative orientation of the nucleic acid sequences is such that in the encoded fusion protein, the N-terminal of the β_2m protein is linked to the C-terminal of the second protein domain. In all instances, the various nucleic acid sequences that comprise the β_2m fusion protein construct (e.g., s, l, x and y) are operably linked such that the elements are situated in a single reading frame.

Nucleic acid constructs expressing fusion proteins may also include regulatory elements
15 such as promoters, enhancers and 3' regulatory regions, the selection of which will be determined based upon the type of cell in which the protein is to be expressed. The constructs are introduced into a vector suitable for expressing the β_2m fusion protein in the selected cell type.

A selected β_2m fusion protein may be obtained by expression in a prokaryotic or eukaryotic expression system, many of which are well known in the art. Heterologous proteins can be produced
20 in prokaryotic cells by placing a strong, regulated promoter and an efficient ribosome binding site upstream of the β_2m fusion protein construct. Suitable promoter sequences include the beta-lactamase, tryptophan (trp), and lambda derived P_L promoters. Prokaryotic expression vectors and expression systems suitable for producing high levels of protein bacterial cells are available commercially and include the pBAD, P_L and Superlinker expression systems produced by Invitrogen
25 (Carlsbad, CA) and the pMAL expression system produced by New England Biolabs (Beverly, MA). In addition, methods and plasmid vectors for producing heterologous proteins in bacteria are described in Sambrook et al. (*Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, New York, 1989). Often, proteins expressed at high levels are found in insoluble inclusion bodies; methods for extracting proteins from these aggregates are described by Sambrook et al. (*Molecular*
30 *Cloning: A Laboratory Manual*, Cold Spring Harbor, New York, 1989, ch. 17). Vectors suitable for the production of intact native proteins include pKC30 (Shimatake and Rosenberg, 1981, *Nature* 292:128), pKK177-3 (Amann and Brosius, 1985, *Gene* 40:183) and pET-3 (Studier and Moffatt, 1986, *J. Mol. Biol.* 189:113). Suitable prokaryotic cells for expression of large amounts of β_2m fusion proteins include *Escherichia coli* and *Bacillus subtilis*.

35 Eukaryotic cells such as Chinese Hamster ovary (CHO), monkey kidney (COS), HeLa, *Spodoptera frugiperda*, and *Saccharomyces cerevisiae* may also be used to express β_2m fusion proteins. Regulatory regions suitable for use in these cells include, for mammalian cells, viral

promoters such as those from CMV, adenovirus and SV40, and for yeast cells, the promoter for 3-phosphoglycerate kinase and alcohol dehydrogenase. Eukaryotic cell expression systems are also commercially available, and include *Pichia pastoris*, *Drosophila*, Baculovirus and Sindbis expression systems produced by Invitrogen (Carlsbad, CA).

5 The transfer of DNA into eukaryotic, in particular human or other mammalian cells, is now a conventional technique. The vectors are introduced into the recipient cells as pure DNA (transfection) by, for example, precipitation with calcium phosphate (Graham and vander Eb, 1973, *Virology* 52:466) or strontium phosphate (Brash et al., 1987, *Mol. Cell Biol.* 7:2013), electroporation (Neumann et al., 1982, *EMBO J* 1:841), lipofection (Felgner et al., 1987, *Proc. Natl. Acad. Sci USA* 84:7413), DEAE dextran (McCuthan et al., 1968, *J. Natl Cancer Inst.* 41:351), microinjection
10 (Mueller et al., 1978, *Cell* 15:579), protoplast fusion (Schafner, 1980, *Proc. Natl. Acad. Sci. USA* 77:2163-7), or pellet guns (Klein et al., 1987, *Nature* 327:70). Alternatively, the nucleic acid molecules can be introduced by infection with virus vectors. Systems are developed that use, for example, retroviruses (Bernstein et al., 1985, *Gen. Engr'g* 7:235), adenoviruses (Ahmad et al., 1986,
15 *J. Virol.* 57:267), or Herpes virus (Spaete et al., 1982, *Cell* 30:295).

 The β_2m fusion protein produced in mammalian cells may be extracted following release of the protein into the supernatant and may be purified using an immunoaffinity column prepared using anti- β_2m antibodies. Alternatively, the β_2m fusion protein may be expressed as a chimeric protein with, for example, β -globin. Antibody to β -globin is thereafter used to purify the chimeric protein.
20 Corresponding protease cleavage sites engineered between the β -globin gene and the nucleic acid sequence encoding the β_2m fusion protein are then used to separate the two polypeptide fragments from one another after translation. One useful expression vector for generating β -globin chimeric proteins is pSG5 (Stratagene).

 By way of example cDNA encoding a β_2m fusion protein with an N-terminal methionine to
25 initiate translation may be subcloned into pET21-d (Novagen) which directs recombinant protein to inclusion bodies. Alternatively, commercially available insect cell expression systems such as the Baculovirus Expression Vector System from Pharmingen (San Diego, CA) can be used for the combined expression and folding of β_2m and β_2m fusion proteins, where the expressed proteins will require only subsequent purification.

30 Proteins expressed as described above may be further purified by immunoaffinity column and used directly to treat mammalian cells. Alternatively, expression of a β_2m fusion protein in a mammalian cell may be obtained by introducing a vector carrying a nucleic acid sequence encoding the protein into the cell. Nucleic acid molecules encoding β_2m fusion proteins carrying a signal sequence, such as the β_2m signal sequence are particularly preferred for this purpose.

35 In one aspect of the invention, the β_2m fusion proteins are used as immunotherapeutics, and the mammalian cell is a tumor cell. In many cancer patients, tumor cells escape immune recognition by downregulating MHC and/or co-stimulatory molecule expression. Accordingly, one method of

treatment previously proposed is to remove tumor cells from a patient, introduce into the cells a co-stimulatory molecule such as B7 and then return the cells to the patient (see for example European Patent Application publication number EP 0 733 373 A2). Applying the discovery disclosed herein to those methods, introducing a β_2m fusion protein into tumor cells is expected to provide
5 considerable benefit.

Obtaining β_2m fusion protein expression on the surface of tumor cells may be achieved either by directly incubating tumor cells with a purified preparation of the fusion protein, or by introducing into the cells a vector that expresses the fusion protein. All types of tumor are potentially amenable to treatment by this approach including, for example, carcinoma of the breast, lung,
10 pancreas, ovary, kidney, colon and bladder, as well as melanomas and sarcomas.

Incorporation of the β_2m fusion protein onto the surface of tumor cells by incubation of the purified fusion protein with the cells, may be achieved by incubation of the tumor cells with the recombinant protein (1-5 μM) in serum-free medium for 2-16 hours. Additionally, tumor cells can be treated briefly with a low pH buffer (pH 2.5 to 3.5) to strip endogenous β_2m and peptide from
15 cell-surface MHC I molecules followed by reconstitution with the β_2m fusion protein and relevant MHC I binding peptides for 1-16 hours.

Where a nucleic acid encoding the fusion protein is to be introduced into the tumor cell, the nucleic acid is preferably incorporated into a suitable expression vector. Suitable vectors include plasmid, cosmid and viral vectors, such as retroviruses, adenoviruses and herpesviruses. Disabled
20 viruses, such as those described in U.S. Patent No. 5,665,362 may be employed for this purpose. Because of the high efficiency with which viral vectors infect mammalian cells, viral vectors are expected to offer advantages over other vector types. The vector is then introduced into the tumor cell by one of a range of techniques, such as electroporation, lipofection, co-cultivation with virus-producing cells, or other standard means. In a preferred embodiment, the tumor cells are cells
25 removed from the patient to be treated, but the tumor cells may alternatively be cells from a tumor cell line, such as the human tumor cell lines available from the American Type Culture Collection (ATCC, Manassas, VA). If it is desired to screen the cells to select those into which the vector was introduced, this may be achieved by a number of means, including selecting for expression of the selectable marker if one is used, or screening for expression of the β_2m fusion protein on the surface
30 of the cells. This latter procedure may be conveniently performed using a fluorescence activated cell sorter (FACS).

The tumor cells are subsequently administered to the patient in combination with a suitable carrier such as buffered water, saline, or glycine. In a preferred embodiment, where the tumor cells are cells originally removed from the patient, they are attenuated before being administered to the
35 patient. An attenuated cell is one which is metabolically active but which is no longer able to proliferate. Methods for attenuating tumor cells are well known and include those described in EP 0 733 373 A2.

In an alternative embodiment, cell membranes from the tumor cells, which include the β_2m fusion protein, may be administered to the patient instead of intact tumor cells. A cell membrane preparation can readily be prepared by disrupting or lysing the cells using standard techniques, such as a French Press, freeze-thawing, or sonication. Following disruption of the cells, a membrane enriched fraction may be obtained by centrifugation.

The present invention also encompasses other immunotherapy methods for treating conditions such as cancer, including adoptive immunotherapy. As is known in the art, adoptive immunotherapy involves obtaining lymphoid cells exposed to a particular antigen, culturing those cells *ex vivo* under conditions whereby the activity of the cells is enhanced, and then administering the cells to an individual. The lymphoid cells are preferably T-cells removed from a cancer patient, for example T-cells from a draining lymph node. These T-cells are incubated with tumor cells removed from the patient which have been treated as described above so as to present a β_2m fusion protein on their cell surface. Accordingly, one aspect of the present invention is a form of adoptive immunotherapy in which the incubation of lymphoid cells *ex vivo* is performed in a medium containing tumor cells presenting a β_2m fusion protein prior to administration of the cells to a patient. The technical details of methods for obtaining lymphoid cells, *ex vivo* cultivation of such cells with immune stimulants, and administration to patients are known in the field and are described, for example in U.S. Patent Nos. 4,690,915 ("Adoptive immunotherapy as a treatment modality in humans"), 5,229,115 ("Adoptive immunotherapy with interleukin-7"), 5,631,006 ("Immunotherapy protocol of culturing leukocytes in the presence of interleukin-2 in a hollow fiber cartridge"), and 4,902,288 ("Implantable immunotherapy system using stimulated cells"), and references cited therein.

Production and use of h β_2m S55V and β_2m fusion proteins in vaccine preparations

Methods for the production and use of β_2m as a vaccine adjuvant are known in the art and include those described in U.S. patent No. 5,733,550, which is incorporated herein by reference. Such methods may be applied for the use of h β_2m S55V as well as β_2m fusion proteins in vaccine preparations as well as methods of vaccination.

β_2m fusion proteins may be produced as described above. Nucleic acid molecules encoding forms of h β_2m carrying the S55V amino acid substitution may be produced using standard mutagenesis techniques, such as site directed mutagenesis, or by PCR as described in Example 4 below. The encoded h β_2m S55V may be expressed in and purified from prokaryotic or eukaryotic expression systems, as described above for the β_2m fusion proteins.

β_2m fusion proteins and h β_2m S55V may be used as adjuvants in vaccine preparations, in which case they may be combined with an antigen in a vaccine preparation, or they may be administered to a patient either shortly before or after administration of a conventional vaccine preparation. Preferably, the β_2m fusion protein or h β_2m S55V is administered at the same location

and contemporaneously with the antigen preparation. Where a $h\beta_2m$ fusion protein is used as a vaccine adjuvant, the $h\beta_2m$ component of the protein may be a S55V form of $h\beta_2m$. The protein to which the β_2m is fused will preferably be a co-stimulatory protein such as B7.

Typically, the antigen administered to a patient in conjunction with a β_2m preparation of the present invention (i.e., a preparation of a β_2m fusion molecule or a S55V $h\beta_2m$) will be a peptide antigen that can bind to class I MHC molecules of the patient. Peptide antigens that may be employed include tumor antigens, as well as antigens from pathogenic organisms, including viruses and bacteria. Examples of suitable antigens include HIV gp120, sub-units of influenza nucleoprotein or hemagglutinin, and tumor antigens as discussed by Boon et al., (*Ann. Rev. Immunol.* 12:337-65, 1994); Finn (*Curr. Opin. Immunol.* 5:701-8, 1993) and Sligluff et al. (*Curr. Opin. Immunol.* 6:733-40, 1994). Such antigens may be isolated or extracted from an original source (e.g., tumor cells), or may be produced by recombinant means, or may be chemically synthesized. Vaccination may be accomplished by administering a single peptide antigen or a cocktail of antigens derived from one or more antigen sources.

The β_2m that forms the basis of the β_2m fusion protein or β_2m S55V used in the vaccine compositions and methods of the present invention may be any mammalian β_2m . It may be preferable to use a β_2m derived from the same species of mammal as the mammal to be vaccinated so as to reduce the risk of immune response to the administered β_2m preparation. However, since xenogeneic β_2m is typically not inflammatory *in vivo*, this may not be necessary.

Vaccine preparations according to the present invention may be administered by any known means, including intramuscular and intravenous injection. In its simplest form, the β_2m preparation administered to the mammal is administered in conventional dosage form, preferably combined with a pharmaceutical excipient, carrier or diluent. Suitable pharmaceutical carriers may be solids or liquids, and may include buffers, anti-oxidants such as ascorbic acid, other polypeptides or proteins such as serum albumin, carbohydrates, chelating agents and other stabilizers and excipients. Suitable solid carriers include lactose, magnesium stearate, terra alba, sucrose, talc, stearic acid, gelatin, agar, pectin, acacia and cocoa butter. The amount of a solid carrier will vary widely depending on which carrier is selected, but preferably will be from about 25 mg to about 1g per dose of active agent. Suitable liquid carriers include neutral buffered saline, optionally with suitable preservatives, stabilizers and excipients. The carrier or diluent may also include time delay material well known to the art such as, for example, glycerol distearate, either alone or with a wax. The foregoing examples of suitable pharmaceutical carriers are only exemplary and one of skill in the art will recognize that a very wide range of such carriers may be employed. Liposome-based delivery systems may also be employed to deliver β_2m preparations. Liposome-based systems, which may be employed to provide a measured release of the agent over time into the bloodstream, are well known in the art and are exemplified by the systems described in U.S. Patent Nos. 4,356,167 ("Liposome drug delivery systems"), 5,580,575 ("Therapeutic drug delivery systems"), 5,595,756

("Liposomal compositions for enhanced retention of bioactive agents") and 5,188,837 ("Lipospheres for controlled delivery of substances"), and documents cited therein.

The formulation of the β_2m preparation with a pharmaceutical carrier can take many physical forms, but is preferably a sterile liquid suspension or solution, suitable for direct injection.

5 Preferably, the patient will be administered the β_2m preparation in a formulation as described above (i.e. in combination with a pharmaceutical carrier), wherein the formulation includes a clinically effective amount of the agent. In the context of vaccination, "a clinically effective amount" of the β_2m preparation is an amount sufficient to provide an enhancement of the immune response to the target antigen, i.e., to produce a cytotoxic T-cell response greater than would be presented absent

10 administration of the β_2m preparation. Quantification of the immune response arising from a vaccination may be achieved in any standard way, e.g., lymphoproliferation in response to test antigen *in vitro* or lysis of target cells by specific cytotoxic T-lymphocytes.

It will be appreciated that a clinically effective dose of a β_2m preparation will vary depending upon the actual β_2m being used (e.g., whether it is a β_2m fusion protein or β_2m S55V

15 alone), and the characteristics of the patient (age, weight, other medications being taken etc.). Thus, the assessment of a clinically effective dosage will ultimately be decided by a physician, veterinarian, or other health care worker familiar with the patient. Typically, administering a β_2m preparation to a mammal as a component of a vaccination regimen will involve administration of from about 10 ng to 1 g of β_2m preparation per dose, with single dose units of from about 10 mg to 100 mg being

20 commonly used, and specific dosages of up to 1 mg or 10 mg also being within the commonly used range. The amount of antigen included in the vaccine preparation that employs a β_2m adjuvant will typically be the same as would be included in vaccine preparations without the β_2m adjuvant, although greater or lesser amounts of antigen may be employed as clinically appropriate.

Where the β_2m is administered to the mammal in a single preparation with the vaccine

25 antigens, the preparation may be formulated simply by mixing a clinically effective amount of the β_2m with the antigen preparation. Vaccines comprising tumor antigens and a β_2m may be prepared from tumor cells which have been transformed to express the β_2m .

EXAMPLES

30 The following examples serve to illustrate the invention.

EXAMPLE 1

Production of β_2m -B7 fusion protein

The murine B7.2:human β_2m fusion protein (mB7 β_2) was made in multiple steps as

35 follows. The extracellular domain of murine B7.2 (Freeman et al., 1993, *J. Exp. Med.* 178:2185-92, GenBank accession number L25606) was amplified by PCR from the plasmid mB7-2 (from Richard Hodes, NIH) using the following two oligonucleotides:

mB7-2 5' PCR Oligo: AGGGTACCATGGTTTCCGTGGAGACGCAAGC (Seq. I.D. No. 8) and mB7-2 3' PCR Oligo: TCGAATTCATGATGCTAGCCCAATACGTTTGAGGAGATGG (Seq. I.D. No. 9) which have embedded restriction sites for cloning (**bold**): KpnI: GAATTC; NcoI: CCATGG; and BspH1: TGATCA.

- 5 The resulting PCR fragment was cut with NcoI and BspH1, and ligated as an amino terminal extension into an NcoI cut bluescript SK vector containing the signal peptide of h β_2 m (Guessow et al., 1987, *J. Immunol.* 139:3132-8, GenBank accession number: M17986), the c-myc tag EQKLISEEDLN (Zhou et al., 1996, *Mol Immunol.* 33:1127-34, Seq. I.D. No. 15), and full length h β_2 m (plasmid #267). The resulting construct (plasmid #392) was then cut with NheI to linearize it
- 10 5' of the myc sequence. Synthetic oligonucleotides encoding a [gly4ser]₃ spacer were engineered with NheI compatible ends and ligated into the linearized vector to create plasmid #396. Finally, the entire coding sequence of wild-type h β_2 m (without a myc tag) was PCR amplified from a h β_2 m cDNA, with the addition of NheI site 5' and a Not I site 3' of the coding sequence. This product was digested with NheI and NotI, and subcloned into plasmid #396 that had also been digested with NheI
- 15 and NotI to generate plasmid #406. This plasmid contained the signal sequence of h β_2 m, followed by the extracellular domain of mB7.2, a 15 amino acid spacer, then mature h β_2 m. For expression in bacteria, the eukaryotic signal sequence was removed. Thus, plasmid #406 was digested with NcoI and NotI to liberate the fusion protein without the signal peptide present, and this was then subcloned into the bacterial expression vector pET21-d that had been linearized with NcoI and NotI.
- 20 Following transfection of the BL21 (DE3) strain of *E. coli*, protein synthesis is induced with IPTG, cells were harvested, lysed, and inclusion bodies washed and solubilized. Following refolding of the recombinant material, it was further purified by gel filtration and/or affinity chromatography with anti- β_2 m antibodies. Experiments in which acid-stripped cells expressing only HLA-A2 were incubated under various conditions with the B7- β_2 m fusion protein produced as described above, and
- 25 then subjected to FACS analysis using conformationally-sensitive antibodies of varying specificities confirmed the following: (1) that the B7 domain of the fusion protein is natively folded; (2) that the β_2 m domain of the fusion protein is natively folded; and (3) that the β_2 m domain of the fusion protein functions to stabilize MHC I expression (data not shown).

30

EXAMPLE 2

The B7- β_2 m fusion protein co-stimulates T-Cells

- The ability of B7- β_2 m to co-stimulate splenic T-cells was determined *in vitro*. Antibodies specific for the B7 receptor CD28 were used as a control. β_2 m, recombinant B7- β_2 m, or anti-CD28 was added to the wells of a microtiter plate and incubated at 37°C for 2 hours to promote binding.
- 35 Thereafter the plates were washed to remove excess reagent, and splenic T-cells from BALB/c mice were added to the wells in the presence of sub-optimal concentrations of the soluble anti-T-cell receptor 2C11. The plates were incubated for 48 hours and then ³H-thymidine was added and

allowed to incorporate into the proliferating cells for 20 hours. At the end of the time period, the cells were removed and the amount of ^3H -thymidine taken up was measured.

The results of these experiments showed that while no T-cell proliferation was observed in wells containing $\beta_2\text{m}$ alone, recombinant B7- $\beta_2\text{m}$ provides co-stimulation to the T-cells at least as effectively as the anti-CD28 antibody. A typical result is shown in FIG. 5.

EXAMPLE 3

Treatment of tumor cells with the $\beta_2\text{m}$ -B7 fusion protein boosts the generation of tumor-specific cytotoxic T-cells

The ability of the B7- $\beta_2\text{m}$ fusion protein to stimulate T-cell recognition and response against tumor cell antigens was compared to the corresponding activity of h $\beta_2\text{m}$ alone. DBA/2 mice were each vaccinated with 3×10^6 syngeneic P815 tumor cells that had been previously incubated in serum-free Iscove's Modified Dulbecco's Medium (SF IMDM) with either $0.2 \mu\text{M}$ B7- $\beta_2\text{m}$, $0.2 \mu\text{M}$ h $\beta_2\text{m}$ alone, or no additional reagent. Samples of each type were analyzed by flow cytometry to show that they stained for B7 or h $\beta_2\text{m}$. These tumor cells were then irradiated with 20,000 Rads and injected into the mice. One week later the mice were reimmunized with identically prepared cells. After an additional week, the mice were sacrificed and their spleen cells were restimulated in culture with irradiated P815 stimulator cells for one week. The resulting cultured T-cells were then assayed for ability to kill untreated P815 tumor cells.

The results, depicted in FIG. 6, show that the spleen cells from the mice immunized with the B7- $\beta_2\text{m}$ fusion protein-treated tumor cells were three fold more effective at killing tumor cells than cells from mice immunized with either the h $\beta_2\text{m}$ treated or untreated tumor cells. These data show that the presence of the fusion protein on the tumor cell surface enhances the immune system response to the tumor antigens present on the tumor cell surface.

EXAMPLE 4

Production of high affinity variant of h $\beta_2\text{m}$

A number of variant forms of h $\beta_2\text{m}$ were created and tested for activity using MHC stabilization, T-lymphocyte lysis and myc-h $\beta_2\text{m}$ binding and inhibition assays. The hydrophilic serine 55 (S55) residue of h $\beta_2\text{m}$ that is buried at the h $\beta_2\text{m}$ /heavy chain interface and is situated directly adjacent to an ordered water molecule was identified as the target residue for mutagenesis. This residue was mutagenized to hydrophobic residues of increasing mass (valine, isoleucine, phenylalanine) in order to promote hydrophobic interactions and exclude the ordered water.

The h $\beta_2\text{m}$ sequence variants were constructed by mutating h $\beta_2\text{m}$ cDNA in Bluescript SK (Stratagene, La Jolla, CA) using the ExSite mutagenesis system (Stratagene) according to the manufacturer's protocol. The mutated cDNAs were subcloned into the bacterial expression vector pET-21d(+) (Novagen, Madison, WI) using engineered NcoI and BamHI sites at the 5' and 3' end of

the mature protein sequence, respectively. Oligonucleotides used to create variants of the h β_2 m sequence were as follows:

Sense S55F: 5' TTC TTC AGC AAG GAC TGG TCT TTC 3' (Seq. I.D. No. 4)

Sense S55I: 5' ATT TTC AGC AAG GAC TGG TCT TTC 3' (Seq. I.D. No. 5)

5 Sense S55V: 5' GTG TTC AGC AAG GAC TGG TCT TTC 3' (Seq. I.D. No. 6)

Common antisense: 5' TAA GTC TGA ATG CTC CAC TTT TTC 3' (Seq. I.D. No. 7)

Expression and purification of the mutated h β_2 ms was performed as previously described by Shields et al. (*J. Immunol.* 160:2297-307, 1998). Briefly, h β_2 m constructs in pET-21d(+) were transformed into the BL21(DE3) strain of *E. coli*. At an O.D._{600nm} of 0.6, cultures were induced with
 10 1 mM IPTG for four hours, and inclusion bodies isolated by centrifugation after sonication of bacteria in 200 mM Tris, 2 mM EDTA, 10% Triton X-100, pH 7.6 and washing in 200 mM Tris, 2 mM EDTA, pH 7.6. Inclusion bodies were solubilized in 6 M Guanidine-HCl containing 0.3 M DTT, 100 mM Tris, pH 8.0, and a mixture of protease inhibitors (5 μ g/ml Leupeptin, 0.5 mM AEBSF, 1% Aprotinin). Following overnight dialysis in 6 M Guanidine pH 2.0, recombinant protein
 15 was refolded over 72 h in 0.4 M Arginine, 5 mM oxidized glutathione, 100 mM Tris, 2 mM EDTA at 10°C. Following refolding, preparations were dialyzed exhaustively against 0.4 M Arginine, 100 mM Tris, 2 mM EDTA, pH 8.0 and then PBS at 4°C. Preparations were purified as a single peak by preparative FPLC on a Superdex 75pg gel filtration column (Pharmacia, Uppsala, Sweden), concentrated using Centriprep-3 concentrating units (Amicon, Beverly, MA), sterile filtered and
 20 concentrations calculated based on O.D._{280nm} readings. Recombinant h β_2 m was judged to be \geq 95% pure based on analysis by SDS-PAGE, and analytical FPLC.

EXAMPLE 5

h β_2 m S55V produces enhanced MHC I stabilization

25 An HLA stabilization assay was used to screen the h β_2 m variants. A number of HLA alleles, HLA-A1, HLA-A2, and HLA-A3, were analyzed in order to determine whether any of the point mutants exhibited allele-specific effects.

In this and the following experiments, cell lines, monoclonal antibodies (mAbs) and peptides used were as follows:

30 **Cell Lines and antibodies:** Hmy2.C1R cells (Storkus et al., 1987, *J. Immunol.* 138:1657) were stably transfected with HLA-A1, -A2, and -A3 as previously described (Winter et al., 1991, *J. Immunol.* 146:3508; DiBrino et al., 1993, *J. Immunol.* 151:5930). HLA-A2/HTLV-1 TAX 11-19 peptide-specific CTL clone N1218 and HLA-A3/Influenza NP 265-273 peptide clone 2G12 were isolated and restimulated as previously described by Biddison et al. (*J. Immunol.* 159:2018, 1997).
 35 All mAbs were used as culture supernatants grown in DMEM supplemented with 10% fetal calf serum, 20 mM HEPES, 2 mM L-glutamine, 1% non-essential amino acids, 1% Pen-strep, and 0.04 mg/ml of Gentamicin sulfate (complete DMEM). GAP.A3 (HLA-A3 specific), and BB7.5 (pan-

HLA-ABC specific) hybridomas were obtained from the American Type Culture Collection (Manassas, VA). The myc-specific 9E10 hybridoma has been previously described by Evan et al. (*Mol Cell Biol.* 5:3610-6, 1985). Unless otherwise noted all solutions used for cell growth were obtained from Biofluids (Rockville, MD).

5 **Peptides:** The peptides used were the HLA-A1 binding ornithine decarboxylase 309-317 (OD 309): SSEQTFMY (Seq. I.D. No. 16); the HLA-A2 binding HTLV-1 TAX 11-19: LLFGYPVYV (Seq. I.D. No. 17) and HIV gag 77-85: SLYNTVATL (Seq. I.D. No. 18); and the HLA-A3 binding pn2a.A3: KLYEKVYTYK (Seq. I.D. No. 19) and influenza NP 265-273: ILRGSVAHK (Seq. I.D. No. 20) (DiBrino et al., 1993, *Proc. Natl. Acad. Sci USA* 90:1508; DiBrino et al., 1994, *J. Immunol.* 152:620; Honma et al., 1997, *J. Neuroimmunol.* 73:7; Parker et al., 1992, *J. Immunol.* 149:3580; Parker et al., 1994, *J. Immunol.* 152:163; and Parker et al., 1995, *Immunol. Res.* 14:34). These peptides were purchased from Bachem (Torrance, CA) or provided by Dr. John E. Coligan (Natl. Inst. of Allergy and Infectious Diseases, NIH). All peptides were purified by reverse phase HPLC and were >95% pure as determined by analytical HPLC and mass spectrometry.

15 The MHC stabilization was done essentially as described previously (Bremers et al., 1995, *J. Immunol. Emphasis Tumor Immunol.* 18:77; van der Burg et al., 1995, *Hum. Immunol.* 44:189; Sugawara et al., 1987, *J. Immunol. Methods* 100:83) with minor modifications. Briefly, Hmy2.C1R cells (Storkus et al., 1987, *J. Immunol.* 138:1657) were stably transfected with HLA-A1, -A2, and -A3 as previously described (Winter et al., 1991; *J. Immunol.* 146:3508; DiBrino et al., 1993, *J. Immunol.* 151:5930). Hmy2.C1R-A1, -A2, and -A3 cells were washed twice with PBS, resuspended in 0.13 M citric acid, 66 mM Na₂HPO₄, pH 2.9 (pH 3.2 for A2 cells), for 90 seconds at 4°C, washed with two 50 ml changes of IMDM, and resuspended in SF IMDM (identical to SF DMEM using IMDM instead). 10⁵ cells per well were added to a 96 well microtiter plate containing hybridoma supernatants, peptide, and hβ₂m dilutions in a total volume of 150 μl. HLA-A1 transfected
25 Hmy2.C1R cells were combined with BB7.5 mAb and 10 μg/ml A1-binding OD 309 peptide. HLA-A2 transfected Hmy2.C1R cells were combined with BB7.5 mAb and 2.5 μg/ml A2-binding HIV gag peptide. HLA-A3 transfected Hmy2.C1R cells were combined with GAP.A3 mAb and 1.25 μg/ml A3-binding pn2a.A3 peptide. After a four hour incubation at 23°C, cells were washed twice with FACS buffer (PBS, 2 mg/ml BSA, 0.02% NaN₃) and stained with FITC-conjugated goat anti-mouse IgG (H+L) F(ab')₂ fragment (Cappel/Organon Teknika, Durham, NC) for one hour at 4°C.
30 Cells were washed twice with FACS buffer and fixed in 1% formaldehyde in PBS followed by flow cytometric analysis on a FACScan II machine (Becton Dickinson, Mountain View, CA).

Figure 7 demonstrates the ability of the S55 variant hβ₂m to stabilize HLA-A1 (FIG. 7a), HLA-A2 (FIG. 7b), and HLA-A3 (FIG. 7c) in the presence of a specific binding peptide and an
35 appropriate HLA-specific antibody. The S55V variant ("X" symbol in FIG. 7) stabilized HLA-A1 and HLA-A3 approximately 2-fold and 3-fold better respectively, than wild-type hβ₂m (diamonds) at a molar level, and effects on HLA-A2 stabilization by S55V were slightly better than those observed

with wild-type $h\beta_2m$. S55F (squares) was similar to wild-type $h\beta_2m$ for all alleles tested, while the effects of S55I (triangles) varied depending on the allele (better with HLA-A1 and worse with -A2 and -A3).

5

EXAMPLE 6 **$h\beta_2m$ S55V binds to MHC I with a higher affinity than wild-type $h\beta_2m$**

The antibodies used in the experiments in the previous Example were selected due to their dependence on both $h\beta_2m$ and peptide in order to detect "complete" molecules, i.e. heavy chain/ $h\beta_2m$ /peptide natively folded trimeric complexes. Since this binding assay requires the presence of an antibody in addition to $h\beta_2m$ and peptide (van der Burg, 1995, *Hum. Immunol.* 44:189), there was the possibility that the antibody itself exerted an effect that is specific for a particular $h\beta_2m$ mutant. Due to concerns regarding the potential contribution of the antibodies to the stabilization of cell-surface MHC I complexes, a binding inhibition assay was developed that directly measures the relative abilities of $h\beta_2m$ s to bind to MHC I molecules. This assay format requires a labeled $h\beta_2m$ to measure the inhibition. Methods of labeling such as biotinylation and iodination are random reactions and create multiply labeled species needing further purification prior to use in a proper competition assay (Hochman et al., 1988, *J. Immunol.* 140:2322). However, endogenous labeling with an epitope tag creates an uniquely labeled species of $h\beta_2m$. Additionally, tyrosine and lysine residues (common targets of biotinylation and iodination) known to be at the MHC heavy chain/ $h\beta_2m$ interface would not be affected with an endogenous label. Therefore an epitope tag (myc) was engineered onto the amino terminus of $h\beta_2m$ and the ability of the various $h\beta_2m$ mutants to compete with the myc- $h\beta_2m$ for cell-surface binding using the anti-myc mAb 9E10 was studied.

To establish the functional activity of the myc- $h\beta_2m$ itself, direct binding studies were performed. Briefly, Hmy2.C1R transfectant cells at 2.5×10^5 per tube in a 500 μ l volume were incubated at 37°C for 16 hours in SF IMDM with 2.5 μ M myc- β_2m , 20 μ g/ml peptide and the indicated concentrations of inhibitor β_2m . The OD 309 peptide was used for HLA-A1, the HIV gag peptide for HLA-A2, and the pn2a.A3 peptide for HLA-A3. Cells were washed three times in plain IMDM followed by incubation with 9E10 (anti-myc) hybridoma supernatant at 4°C for one hour. After washing with IMDM, cells were stained for one hour with FITC anti-mouse IgG at 4°. Cells were washed a final time in FACS buffer and analyzed by flow cytometry, gating on live (propidium iodide-excluding) cells. In the presence of an appropriate peptide there was concentration-dependent myc- β_2m binding for all alleles studied. However, when cells were incubated with myc- β_2m in the absence of peptide, no appreciable myc- $h\beta_2m$ binding was observed.

The relative abilities of wild-type $h\beta_2m$ and S55V to inhibit the binding of myc- $h\beta_2m$ to HLA molecules were next compared using an inhibition assay. The inhibition assay was identical to the binding assay with the following modifications: 2.5 μ M myc- $h\beta_2m$ was used in all cases and

different concentrations of non-myc labeled recombinant h β_2 m were included to inhibit myc-h β_2 m binding to cell surface MHC molecules. Percent inhibition was calculated by the following equation: $(1 - ((\text{experimental-background}) / (\text{no inhibitor-background}))) \times 100$. 10-20,000 gated events per sample were counted, and all experiments were repeated at least twice. Compared with wild-type h β_2 m, the S55V mutant inhibited myc-h β_2 m binding about 2.5-fold better at a molar level for HLA-A1, -A2, and -A3 (FIG. 8). These results demonstrate the higher relative affinity of the S55V mutant compared to wild-type h β_2 m for HLA-A1 (FIG. 8a), -A2 (FIG. 8b) and -A3 (FIG. 8c).

EXAMPLE 7

h β_2 m S55V enhances CTL recognition of target cells

The effectiveness of h β_2 m in facilitating exogenous peptide loading of MHC I molecules was measured using a CTL lysis assay (Depierreux et al., 1997, *J. Immunol. Methods* 203:77) as follows: Hmy2.C1R cells transfected with HLA-A2 or HLA-A3 were resuspended to 4×10^6 cells/ml in complete DMEM supplemented with 20 μ M BATD (which forms a fluorescent chelate with Europium; Wallac, Gaithersburg, MD) and incubated at 37°C for 30-60 minutes. Cells were resuspended in 10 ml of serum free (SF) CTL medium (IMDM supplemented with 5 mg/ml bovine serum albumin (Sigma, St. Louis, MO), 2 mM L-glutamine, 1.25 mM sulfinpyrazone (Sigma), and 1% Pen-strep), centrifuged and washed once more with SF CTL medium. Cells were then pulsed with peptide and/or β_2 m in SF CTL medium for 60-90 minutes at 37°C. Cells were washed twice in SF CTL medium, resuspended in CTL medium (5% fetal calf serum in lieu of BSA) and combined at the designated effector:target ratio with CTL clones in round bottom microtiter plates (CTL clone N1218 at an E:T ratio of 4:1 (FIG. 9a) or the NP-specific HLA-A3 restricted CTL clone 2711 at an E:T ratio of 2:1 (FIG. 9b)). Plates were gently centrifuged at 100 x g for 2 minutes and then incubated at 37°C for 2 hours. Finally, plates were centrifuged at 300 x g and 20 μ l per well was transferred to 200 μ l of 0.3 M acetic acid, 60 mM sodium acetate, 7.5 μ g/ml Europium (Aldrich, Milwaukee, WI), and the plate read on a Wallac 1234 DELFIA Fluorometer. Percent specific lysis was calculated with the following equation:

$$100 \times ((\text{experimental-blank}) - (\text{spontaneous-blank})) / ((\text{maximum-blank}) - (\text{spontaneous-blank})).$$

In this assay, target cell lysis not only correlates with the loading of a specific peptide antigen, but it also demonstrates that the peptide is bound in an immunologically relevant manner. Hmy2.C1R-A2 target cells (a human lymphoblastoid cell essentially null for HLA molecules except for the transfected HLA-A2.1) (Winter et al., 1991; *J. Immunol.* 146:3508; DiBrino et al., 1993, *J. Immunol.* 151:5930) were pulsed with a suboptimal concentration of HTLV-1 TAX peptide (9.3×10^{-12} M/l) or control A2-binding HIV gag peptide at 1×10^{-9} M/l for 90 minutes in serum-free CTL medium in the absence or presence of increasing concentrations of purified, recombinant h β_2 m and then used as targets in a conventional lysis assay. The presence of wild-type h β_2 m dramatically increased the specific lysis by the TAX-specific CTL clone in a dose-dependent fashion. Using this

suboptimal concentration of peptide there was 20% lysis in the absence of h β_2 m. Addition of 8 μ M h β_2 m increases the lysis to the maximal observed at this E:T ratio. In the absence of h β_2 m 100 fold higher concentration of peptide would be required to achieve comparable levels of lysis (data not shown).

5 Having established the ability of wild-type h β_2 m to enhance the loading of antigenic peptide onto cells, the activity of the S55V variant in this assay was examined. Two CTL clones, specific for an HTLV-1 TAX peptide in the context of HLA-A2 and an influenza nucleoprotein peptide in the context of HLA-A3, were used in the assay described above using Hmy2.C1R transfectants pulsed with a suboptimal concentration of antigenic peptide (NP 265-273, 1×10^{-10} M/l) or control A3-binding pn2a.A3 peptide at 1×10^{-6} M/l. The S55V mutant was 4-fold more effective at a molar level than wild-type h β_2 m at enhancing target cell lysis for HLA-A2 (FIG. 9a) and 6 to 7-fold better for HLA-A3 (FIG. 9b). Controls with irrelevant -A2 and -A3 binding peptides with the highest concentrations of h β_2 m used resulted in only background levels of killing. Additionally, multiple TAX-specific A2-restricted clones displayed similar levels of S55V enhanced killing relative to wild-type β_2 m (data not shown).

20 In view of the many possible embodiments to which the principles of the invention may be applied, it will be recognized that the foregoing examples are offered for purposes of illustration and do not limit the scope of the invention. Rather, the scope of the invention is defined by the following claims. We therefore claim as our invention all that comes within the scope and spirit of these claims.

We claim:

1. A fusion protein comprising a first amino acid sequence and a second amino acid sequence, wherein the second amino acid sequence is a β_2 -microglobulin.
2. A fusion protein according to claim 1 wherein the second amino acid sequence is a human β_2 -microglobulin.
3. A fusion protein according to claim 1 wherein the second amino acid sequence is h β_2 m S55V.
4. A fusion protein comprising first and second domains, wherein the second domain is β_2 m.
5. A fusion protein according to claim 4 wherein the first domain joined to the amino terminal of the second domain.
6. A fusion protein according to claim 4 wherein the second domain is h β_2 m.
7. A fusion protein according to claim 4 wherein the first domain is a co-stimulatory protein.
8. A fusion protein according to claim 7 wherein the co-stimulatory protein is selected from the group consisting of B7.1 and B7.2.
9. A fusion protein according to claim 4 wherein the first domain is an integrin, a cytokine or a cell adhesion molecule.
10. A fusion protein according to claim 6 wherein the h β_2 m is h β_2 m S55V.
11. A fusion protein according to claim 4 wherein the first and second domains are linked by a peptide linker.
12. A fusion protein according to claim 4 wherein the fusion protein further comprises a signal peptide joined to the N terminus of the first domain.
13. A fusion protein according to claim 12 wherein the signal peptide is a β_2 m signal peptide.
14. A fusion protein according to claim 10 wherein the h β_2 m S55V has an amino acid sequence as shown in Seq. I.D. No. 10.
15. A fusion protein according to claim 6 wherein the protein has an amino acid sequence selected from the group consisting of the sequences shown in Seq. I.D. Nos. 2 and 3.
16. A recombinant nucleic acid molecule encoding a protein according to claim 4.
17. A vector comprising a nucleic acid molecule according to claim 16.
18. A transgenic cell comprising a nucleic acid molecule according to claim 16.
19. A cell having a cell membrane comprising a fusion protein according to claim 4.
20. A cell according to claim 19 wherein the cell is a tumor cell.
21. A protein comprising a structure X-Y wherein X is a protein domain and Y is a beta-2 microglobulin.

22. A protein according to claim 21 comprising a structure X-L-Y wherein L is a linker peptide.
23. A protein according to claim 22 comprising S-X-L-Y wherein S is a signal peptide.
- 5 24. A protein according to claim 23 wherein the signal peptide is a β_2m signal peptide.
25. A nucleic acid molecule encoding a protein according to claim 21.
26. A protein according to claim 21 wherein the protein has an amino acid sequence as shown in Seq. I.D. No. 2 or Seq. I. D. No. 3.
- 10 27. A method of enhancing the immune response of a mammal to an antigen presented on the surface of a cell, the method comprising:
- (a) contacting the cell with a fusion protein according to claim 4 such that the fusion protein is presented on the surface of the cell; and
 - (b) administering the cell to a mammal.
- 15 28. The method of claim 27 wherein the first amino acid sequence is a co-stimulatory molecule.
29. The method of claim 28 wherein the co-stimulatory molecule is B7.1 or B7.2.
30. The method of claim 29 wherein the cell is a tumor cell.
31. A method of enhancing the immune response of a mammal to an antigen
- 20 presented on the surface of a cell, the method comprising:
- (a) transforming the cell with a nucleic acid molecule according to claim 16, such that expression of the nucleic acid molecule results in expression of a fusion protein encoded by the nucleic acid molecule being presented on the surface of the cell; and
 - (b) administering the cell to a mammal.
- 25 32. The method of claim 31 wherein the first amino acid sequence is a co-stimulatory molecule.
33. The method of claim 32 wherein the co-stimulatory molecule is B7.1 or B7.2.
34. The method of claim 33 wherein the cell is a tumor cell.
35. A human β_2 -microglobulin molecule having a valine residue at position 55.
- 30 36. A human β_2 -microglobulin molecule according to claim 35, wherein the molecule comprises the amino acid sequence shown in Seq. I.D. No. 10.
37. A vaccine preparation comprising at least one antigen and a molecule selected from the group consisting of
- (a) a human β_2 -microglobulin molecule having a valine at position 55; and
 - (b) a fusion protein comprising a first amino acid sequence and a second
- 35 amino acid sequence, wherein the second amino acid sequence is a β_2 -microglobulin.

38. A vaccine preparation according to claim 37(b) wherein the β_2 -microglobulin is h β_2 m S55V.

39. A vaccine preparation according to claim 37 wherein the antigen is selected from the group consisting of bacterial, viral and tumor antigens.

5 40. A method of vaccinating a mammal, comprising administering to the mammal a vaccine preparation according to claim 37.

41. A method of vaccinating a mammal, comprising administering to the mammal an antigen and a microglobulin protein selected from the group consisting of:

- 10 (a) a human β_2 -microglobulin protein having a valine at position 55; and
(b) a fusion protein comprising a first amino acid sequence and a second amino acid sequence, wherein the second amino acid sequence is a β_2 -microglobulin.

42. A method of stimulating a tumor-reactive cytotoxic T-cell response, comprising:

- 15 (a) isolating T-cells from a patient having a tumor;
(b) isolating tumor cells from the patient;
(c) incubating the tumor cells with a fusion protein according to claim 4, such that the fusion protein is presented on the surface of the tumor cells;
(d) incubating the T-cells in the presence of the fusion protein-presenting tumor cells to increase the number of tumor-reactive T-cells; and
20 (e) administering a therapeutically effective dose of the tumor-reactive T-cells to the patient.

1/7


FIG. 1

M S R S V A L A V L A L L S L S G L E A I Q R T P K I Q V Y 10
1
S R H P A E N G K S N F L N C Y V S G F H P S D I E V D L L 40
20 30
K N G E R I E K V E H S D L S F S K D W S F Y L L Y Y T E F 70
50 60
T P T E K D E Y A C R V N H V T L S Q P K I V K W D R D M 90
80

FIG. 3

M V S V E T Q A Y F N G T A Y L P C P F T K A Q N I S L S E 30
L V V F W Q D Q Q K L V L Y E H Y L G T E K L D S V N A K Y 60
L G R T S F D R N N W T L R L H N V Q I K D M G S Y D C F I 90
Q K K P P T G S I I L Q Q T L T E L S V I A N F S E P E I K 120
L A Q N V T G N S G I N L T C T S K Q G H P K P K K M Y F L 150
I T N S T N E Y G D N M Q I S Q D N V T E L F S I S N S L S 180
L S F P D G V W H M T V V C V L E T E S M K I S S K P L N F 210
T Q E F P S P Q T Y W A S T S G G G G S G G G G S G G G A S 240
I Q R T P K I Q V Y S R H P A E N G K S N F L N C Y V S G F 270
H P S D I E V D L L K N G E R I E K V E H S D L S F S K D W 300
S F Y L L Y Y T E F T P T E K D E Y A C R V N H V T L S Q P 330
K I V K W D R D M 339

FIG. 2



3/7

FIG. 4

M S R S V A L A V L A L L S L S G L E A V S V E T Q A Y F N 30
G T A Y L P C P F T K A Q N I S L S E L V V F W Q D Q Q K L 60
V L Y E H Y L G T E K L D S V N A K Y L G R T S F D R N N W 90
T L R L H N V Q I K D M G S Y D C F I Q K K P P T G S I I L 120
Q Q T L T E L S V I A N F S E P E I K L A Q N V T G N S G I 150
N L T C T S K Q G H P K P K K M Y F L I T N S T N E Y G D N 180
M Q I S Q D N V T E L F S I S N S L S L S F P D G V W H M T 210
V V C V L E T E S M K I S S K P L N F T Q E F P S P Q T Y W 240
A S T S G G G G S G G G G S G G G A S I Q R T P K I Q V Y S 270
R H P A E N G K S N F L N C Y V S G F H P S D I E V D L L K 300
N G E R I E K V E H S D L S F S K D W S F Y L L Y Y T E F T 330
P T E K D E Y A C R V N H V T L S Q P K I V K W D R D M 358

4/7




FIG. 5




FIG. 6

5/7

FIG. 7a




FIG. 7b




FIG. 7c




FIG. 8a




FIG. 8b





FIG. 8c



7/7

FIG. 9a





FIG. 9b



SEQUENCE LISTING

<110> National Institutes of Health

<120> B2 Microglobulin Fusion Proteins and High Affinity Variants

<130> 52590

<140>

<141>

<150> 60/088,813

<151> 1998-06-10

<160> 20

<170> PatentIn Ver. 2.0

<210> 1

<211> 119

<212> PRT

<213> Homo sapiens

<400> 1

Met Ser Arg Ser Val Ala Leu Ala Val Leu Ala Leu Leu Ser Leu Ser
1 5 10 15

Gly Leu Glu Ala Ile Gln Arg Thr Pro Lys Ile Gln Val Tyr Ser Arg
20 25 30

His Pro Ala Glu Asn Gly Lys Ser Asn Phe Leu Asn Cys Tyr Val Ser
35 40 45

Gly Phe His Pro Ser Asp Ile Glu Val Asp Leu Leu Lys Asn Gly Glu
50 55 60

Arg Ile Glu Lys Val Glu His Ser Asp Leu Ser Phe Ser Lys Asp Trp
65 70 75 80

Ser Phe Tyr Leu Leu Tyr Tyr Thr Glu Phe Thr Pro Thr Glu Lys Asp
85 90 95

Glu Tyr Ala Cys Arg Val Asn His Val Thr Leu Ser Gln Pro Lys Ile
100 105 110

Val Lys Trp Asp Arg Asp Met
115

<210> 2

<211> 339

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: fusion protein

<400> 2

Met Val Ser Val Glu Thr Gln Ala Tyr Phe Asn Gly Thr Ala Tyr Leu
1 5 10 15

Pro Cys Pro Phe Thr Lys Ala Gln Asn Ile Ser Leu Ser Glu Leu Val
 20 25 30
 Val Phe Trp Gln Asp Gln Gln Lys Leu Val Leu Tyr Glu His Tyr Leu
 35 40 45
 Gly Thr Glu Lys Leu Asp Ser Val Asn Ala Lys Tyr Leu Gly Arg Thr
 50 55 60
 Ser Phe Asp Arg Asn Asn Trp Thr Leu Arg Leu His Asn Val Gln Ile
 65 70 75 80
 Lys Asp Met Gly Ser Tyr Asp Cys Phe Ile Gln Lys Lys Pro Pro Thr
 85 90 95
 Gly Ser Ile Ile Leu Gln Gln Thr Leu Thr Glu Leu Ser Val Ile Ala
 100 105 110
 Asn Phe Ser Glu Pro Glu Ile Lys Leu Ala Gln Asn Val Thr Gly Asn
 115 120 125
 Ser Gly Ile Asn Leu Thr Cys Thr Ser Lys Gln Gly His Pro Lys Pro
 130 135 140
 Lys Lys Met Tyr Phe Leu Ile Thr Asn Ser Thr Asn Glu Tyr Gly Asp
 145 150 155 160
 Asn Met Gln Ile Ser Gln Asp Asn Val Thr Glu Leu Phe Ser Ile Ser
 165 170 175
 Asn Ser Leu Ser Leu Ser Phe Pro Asp Gly Val Trp His Met Thr Val
 180 185 190
 Val Cys Val Leu Glu Thr Glu Ser Met Lys Ile Ser Ser Lys Pro Leu
 195 200 205
 Asn Phe Thr Gln Glu Phe Pro Ser Pro Gln Thr Tyr Trp Ala Ser Thr
 210 215 220
 Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Ala Ser
 225 230 235 240
 Ile Gln Arg Thr Pro Lys Ile Gln Val Tyr Ser Arg His Pro Ala Glu
 245 250 255
 Asn Gly Lys Ser Asn Phe Leu Asn Cys Tyr Val Ser Gly Phe His Pro
 260 265 270
 Ser Asp Ile Glu Val Asp Leu Leu Lys Asn Gly Glu Arg Ile Glu Lys
 275 280 285
 Val Glu His Ser Asp Leu Ser Phe Ser Lys Asp Trp Ser Phe Tyr Leu
 290 295 300
 Leu Tyr Tyr Thr Glu Phe Thr Pro Thr Glu Lys Asp Glu Tyr Ala Cys
 305 310 315 320
 Arg Val Asn His Val Thr Leu Ser Gln Pro Lys Ile Val Lys Trp Asp
 325 330 335

Arg Asp Met

<210> 3

<211> 358

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: fusion protein

<400> 3

Met	Ser	Arg	Ser	Val	Ala	Leu	Ala	Val	Leu	Ala	Leu	Leu	Ser	Leu	Ser
1				5					10					15	

Gly	Leu	Glu	Ala	Val	Ser	Val	Glu	Thr	Gln	Ala	Tyr	Phe	Asn	Gly	Thr
			20					25					30		

Ala	Tyr	Leu	Pro	Cys	Pro	Phe	Thr	Lys	Ala	Gln	Asn	Ile	Ser	Leu	Ser
		35					40					45			

Glu	Leu	Val	Val	Phe	Trp	Gln	Asp	Gln	Gln	Lys	Leu	Val	Leu	Tyr	Glu
	50					55					60				

His	Tyr	Leu	Gly	Thr	Glu	Lys	Leu	Asp	Ser	Val	Asn	Ala	Lys	Tyr	Leu
65					70					75					80

Gly	Arg	Thr	Ser	Phe	Asp	Arg	Asn	Asn	Trp	Thr	Leu	Arg	Leu	His	Asn
				85					90					95	

Val	Gln	Ile	Lys	Asp	Met	Gly	Ser	Tyr	Asp	Cys	Phe	Ile	Gln	Lys	Lys
			100					105					110		

Pro	Pro	Thr	Gly	Ser	Ile	Ile	Leu	Gln	Gln	Thr	Leu	Thr	Glu	Leu	Ser
		115					120					125			

Val	Ile	Ala	Asn	Phe	Ser	Glu	Pro	Glu	Ile	Lys	Leu	Ala	Gln	Asn	Val
	130					135					140				

Thr	Gly	Asn	Ser	Gly	Ile	Asn	Leu	Thr	Cys	Thr	Ser	Lys	Gln	Gly	His
145					150					155					160

Pro	Lys	Pro	Lys	Lys	Met	Tyr	Phe	Leu	Ile	Thr	Asn	Ser	Thr	Asn	Glu
				165					170					175	

Tyr	Gly	Asp	Asn	Met	Gln	Ile	Ser	Gln	Asp	Asn	Val	Thr	Glu	Leu	Phe
			180					185					190		

Ser	Ile	Ser	Asn	Ser	Leu	Ser	Leu	Ser	Phe	Pro	Asp	Gly	Val	Trp	His
		195					200					205			

Met	Thr	Val	Val	Cys	Val	Leu	Glu	Thr	Glu	Ser	Met	Lys	Ile	Ser	Ser
	210					215					220				

Lys	Pro	Leu	Asn	Phe	Thr	Gln	Glu	Phe	Pro	Ser	Pro	Gln	Thr	Tyr	Trp
225					230					235					240

Ala	Ser	Thr	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly
				245					250					255	

Gly Ala Ser Ile Gln Arg Thr Pro Lys Ile Gln Val Tyr Ser Arg His
 260 265 270
 Pro Ala Glu Asn Gly Lys Ser Asn Phe Leu Asn Cys Tyr Val Ser Gly
 275 280 285
 Phe His Pro Ser Asp Ile Glu Val Asp Leu Leu Lys Asn Gly Glu Arg
 290 295 300
 Ile Glu Lys Val Glu His Ser Asp Leu Ser Phe Ser Lys Asp Trp Ser
 305 310 315 320
 Phe Tyr Leu Leu Tyr Tyr Thr Glu Phe Thr Pro Thr Glu Lys Asp Glu
 325 330 335
 Tyr Ala Cys Arg Val Asn His Val Thr Leu Ser Gln Pro Lys Ile Val
 340 345 350
 Lys Trp Asp Arg Asp Met
 355

<210> 4
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 4
 ttcttcagca aggactgggc tttc 24

<210> 5
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 5
 attttcagca aggactgggc tttc 24

<210> 6
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 6
 gtgttcagca aggactgggc tttc 24

<210> 7
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 7

taagtctgaa tgctccactt tttc

24

<210> 8

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 8

agggtagcat ggtttccgtg gagacgcaag c

31

<210> 9

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 9

tcgaattcat gatgctagcc caatacgttt gaggagatgg

40

<210> 10

<211> 99

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Modified hB2m
S55V

<400> 10

Ile	Gln	Arg	Thr	Pro	Lys	Ile	Gln	Val	Tyr	Ser	Arg	His	Pro	Ala	Glu
1				5					10					15	

Asn	Gly	Lys	Ser	Asn	Phe	Leu	Asn	Cys	Tyr	Val	Ser	Gly	Phe	His	Pro
		20					25						30		

Ser	Asp	Ile	Glu	Val	Asp	Leu	Leu	Lys	Asn	Gly	Glu	Arg	Ile	Glu	Lys
		35					40					45			

Val	Glu	His	Ser	Asp	Leu	Val	Phe	Ser	Lys	Asp	Trp	Ser	Phe	Tyr	Leu
	50					55					60				

Leu	Tyr	Tyr	Thr	Glu	Phe	Thr	Pro	Thr	Glu	Lys	Asp	Glu	Tyr	Ala	Cys
65					70					75					80

Arg	Val	Asn	His	Val	Thr	Leu	Ser	Gln	Pro	Lys	Ile	Val	Lys	Trp	Asp
				85					90					95	

Arg Asp Met

<210> 11

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: linker that
can be used in fusion proteins

<400> 11

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
1 5 10 15

<210> 12

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: linker that
can be used in fusion proteins

<400> 12

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Ala Ser
1 5 10 15

<210> 13

<211> 21

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: signal peptide

<400> 13

Lys Tyr Leu Leu Pro Thr Ala Ala Ala Gly Leu Leu Leu Leu Ala Ala
1 5 10 15

Gln Pro Ala Met Ala

20

<210> 14

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: signal peptide

<400> 14

Met Arg Ala Lys Leu Leu Gly Ile Val Leu Thr Pro Ile Ala Ile Ser
1 5 10 15

Phe Ala Ser Thr

20

<210> 15

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: c-myc tag

<400> 15

Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu Asn
1 5 10

<210> 16

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: ornithine
decarboxylase 309-317

<400> 16

Ser Ser Glu Gln Thr Phe Met Tyr Tyr
1 5

<210> 17

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: HTLV TAX 11-19

<400> 17

Leu Leu Phe Gly Tyr Pro Val Tyr Val
1 5

<210> 18

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: HIV gag 77-85

<400> 18

Ser Leu Tyr Asn Thr Val Ala Thr Leu
1 5

<210> 19

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: pn2a.A3

<400> 19

Lys Leu Tyr Glu Lys Val Tyr Thr Tyr Lys
1 5 10

<210> 20

<211> 9

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: influenza NP
265-273

<400> 20

Ile Leu Arg Gly Ser Val Ala His Lys
1 5

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/12309

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/13 C12N15/62 C07K14/705 A61K48/00 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	UGER ET AL.: "Creating CTL targets with epitope-linked beta 2-microglobulin constructs" THE JOURNAL OF IMMUNOLOGY, vol. 160, no. 4, 15 February 1998 (1998-02-15), pages 1598-1605, XP002115504 page 1599 -page 1603; figure 1 ---	1,2,4-6, 11-13, 16-25, 27,30, 31,34, 37,39-42
X	MOTTEZ ET AL.: "Cells expressing a major histocompatibility complex class I molecule with a single covalently bound peptide are highly immunogenic" JOURNAL OF EXPERIMENTAL MEDICINE, vol. 181, no. 2, February 1995 (1995-02), pages 493-502, XP000654243 page 495 -page 501; figures 1,3,7-9; table 2 ---	1,4,5,9, 11,12, 16,18, 19, 21-23, 25,27, 28,37,41 7,8,28, 29,32,33
Y	---	---



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

23 September 1999

Date of mailing of the international search report

06/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

van Klompenburg, W

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/12309

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	GERSTMAYER ET AL: "Costimulation of T cell proliferation by a chimeric B7-2 antibody fusion protein specifically targeted to cells expressing the erbB2 proto-oncogene" THE JOURNAL OF IMMUNOLOGY, vol. 158, no. 10, 15 May 1997 (1997-05-15), pages 4584-4590, XP002116142 cited in the application figures 1,4,5 ----	7,8,28, 29,32,33
X	US 5 574 205 A (KUCHERLAPATI RAJU ET AL) 12 November 1996 (1996-11-12) figure 5; example 5 ----	1,2,4-6, 9,11,12, 16-19, 21,25
X	WO 94 24290 A (BRITISH BIO TECHNOLOGY ;EDWARDS RICHARD MARK (GB); HUNTER MICHAEL) 27 October 1994 (1994-10-27) claims 1-22; examples 5,6,8,9 ----	1,2,4-6, 11,12, 16-19, 21-23, 25,37,41 1-42
A	US 5 529 921 A (LANGLADE-DEMOYEN PIERRE ET AL) 25 June 1996 (1996-06-25) claims 1-12; examples 3,4,5,6D ----	1-42
Y	SHIELDS ET AL: "Characterization of the interactions between mhc class I subunits: a systematic approach for the engineering of higher affinity variants of B2-microglobulin" THE JOURNAL OF IMMUNOLOGY, vol. 160, no. 5, 1 March 1998 (1998-03-01), XP002116153 page 2304 -page 2306; figures 8,9 ----	35-37,41
A	BJORKMAN ET AL.: "Structure of the human class I histocompatibility antigen, HLA-A2" NATURE, vol. 329, 8 October 1987 (1987-10-08), pages 506-512, XP002116341 table 2 ----	10,14, 39,40
Y	US 5 733 550 A (ROCK ET AL.) 31 March 1998 (1998-03-31) cited in the application claims 1-22; figure 1; examples 1-3 ----	35-37,41
A	US 5 733 550 A (ROCK ET AL.) 31 March 1998 (1998-03-31) cited in the application claims 1-22; figure 1; examples 1-3 ----	1-42
	----- -/--	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/12309

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>SHIELDS ET AL.: "The effect of human B2-microglobulin on major histocompatibility complex I peptide loading and the engineering of a high affinity variant"</p> <p>THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 273, no. 43, 23 October 1998 (1998-10-23), pages 28010-28018, XP002115505</p> <p>the whole document</p> <p>-----</p>	35,37, 39-41

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 12309

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
**Remark: Although claims 27-34 and 40-42
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.**
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 15, 26 complete, 1-14, 16-25, 27-34,
37-42 all partially

A fusion protein comprising first and second domains, wherein the second domain is B2-microglobulin.
Said fusion protein wherein the first domain is a co-stimulatory protein. A recombinant DNA molecule encoding said fusion protein.
A method of enhancing the immuneresponse of a mammal to an antigen presented on the surface of a cell comprising contacting the cell with said fusion protein and administering the cell to the mammal.
A method of enhancing the immuneresponse of a mammal to an antigen presented on the surface of a cell comprising transforming the cell with said nucleic acid and administering the cell to the mammal.
Vaccine preparations containing said fusion protein and an antigen. Said vaccine wherein the B-2microglobulin is hB2M S55V. A method of vaccination using said vaccine. A method of stimulating a tumor-reactive cytotoxic T-cell response comprising isolating T-cells and tumor cells from a patient, incubating the tumor cells with said fusion protein, incubating the T-cells with said cells and administering the tumor reactive T-cells to the patient.

2. Claims: 35 ,36 complete and 1-14, 16-25, 27-34,
37-42 partially

A human B2-microglobulin molecule having a valine residue at position 55.
A vaccine preparation comprising at least one antigen and said B2-microglobulin.
A method of vaccinating a mammal comprising administering said vaccine to a mammal.
A fusion protein comprising first and second domains, wherein the second domain is a human B2-microglobulin having a valine residue at position 55.
Said fusion protein wherein the first domain is a co-stimulatory protein. A recombinant DNA molecule encoding said fusion protein.
A method of enhancing the immuneresponse of a mammal to an antigen presented on the surface of a cell comprising contacting the cell with said fusion protein and administering the cell to the mammal.
A method of enhancing the immuneresponse of a mammal to an antigen presented on the surface of a cell comprising transforming the cell with said nucleic acid and administering the cell to the mammal.
Vaccine preparations containing said fusion protein and an

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

antigen. Said vaccine wherein the B-2microglobulin is hB2M S55V. A method of vaccination using said vaccine. A method of stimulating a tumor-reactive cytotoxic T-cell response comprising isolating T-cells and tumor cells from a patient, incubating the tumor cells with said fusion protein, incubating the T-cells with said cells and administering the tumor reactive T-cells to the patient.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/12309

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5574205 A	12-11-1996	US 5413923 A US 5416260 A AU 696372 B AU 1445695 A CA 2180019 A EP 0742723 A JP 9507753 T WO 9517911 A AU 633958 B AU 6150590 A CA 2037907 A EP 0437576 A JP 4501510 T WO 9101140 A	09-05-1995 16-05-1995 10-09-1998 17-07-1995 06-07-1995 20-11-1996 12-08-1997 06-07-1995 11-02-1993 22-02-1991 26-01-1991 24-07-1991 19-03-1992 07-02-1995
WO 9424290 A	27-10-1994	AU 6435294 A AU 6435394 A EP 0692027 A EP 0693125 A WO 9424280 A	08-11-1994 08-11-1994 17-01-1996 24-01-1996 27-10-1994
US 5529921 A	25-06-1996	US 5314813 A US 5827737 A AU 3728193 A EP 0628074 A JP 7504089 T WO 9317095 A	24-05-1994 27-10-1998 13-09-1993 14-12-1994 11-05-1995 02-09-1993
US 5733550 A	31-03-1998	AT 147990 T AU 660278 B AU 7987391 A DE 69124372 D DE 69124372 T DK 531401 T EP 0531401 A ES 2096650 T WO 9116924 A	15-02-1997 22-06-1995 27-11-1991 06-03-1997 07-08-1997 28-07-1997 17-03-1993 16-03-1997 14-11-1991